Reconstructing Supersymmetric High Scale Theories using Precision Data

Werner Porod

University of Zürich

LHC/LC meeting, July 2002

- mSUGRA, GMSB, String
- Procedure
- Numerical Results

G. Blair, W. Porod, P. Zerwas

Phys. Rev. D 63 (2001) 017703; in preparation

Supersymmetry breaking

mSUGRA

Universal input Parameters:

M_0, $M_{1/2}$, A_0, $\tan\beta$, sign(μ)

GMSB

Messenger scale: $M_M = \lambda S$

$\Lambda = F/S$ where $< X >= S + F\theta\theta$

$M_{1/2} \to g(x)\alpha_i\Lambda$,

$M_0^2 \to f(X)\sum C_i\alpha_i^2\Lambda^2$,

$x = \Lambda/M_M$, $f(x), g(x) = (n_5 + 3n_{10})O(1)$

A_0, $\tan\beta$, sign(μ)

String

vevs of dilaton $< s >$ and moduli fields $< t_i >$

give determine the parameters $M_i, M_{0,j}, A_{0,k}$.

In general non-universal boundary conditions.

In addition $\tan\beta$, sign(μ)

Assumptions

1) Real Parameters

2) The generation mixing for sfermions is the same as for fermions:

$M^2_{\tilde{f}ij} = \delta_{ij} M^2_{\tilde{f}i}$,

$a_{f,ij} = \delta_{ij} A_{f,i} Y_{f,i}$

$\Rightarrow$ Each of the three $(\tilde{l}, \tilde{u}, \tilde{d})$ generally 6×6 matrices can be decomposed into three 2×2 matrices

Procedure*

- Input at m_Z: $g_1, g_2, g_3, h_l, h_d, h_u, \tan\beta$ in $\overline{DR}$ scheme
- Using 2-loop RGEs to get the above parameters at the high scale: M_{GUT} (in mSUGRA defined via $g_1 = g_2$) or M_M (in GMSB)
- Evaluating mass parameters down to electroweak scale using 2-loop RGEs†, including threshold effects**
- radiative electroweak symmetry breaking

$$\Rightarrow |\mu|^2 = \frac{m_{H_2}^2 \sin^2\beta - m_{H_1}^2 \cos^2\beta}{\cos 2\beta} - \frac{m_Z^2}{2} + \text{rad.corr.}$$

*for details see e.g. Arason et al., Phys.Rev.D **46** 3945 (1992); W. Porod, **SPHENO**, a program handling **S**upersymmetric **PHENO**menology, hep-ph/0207xxx

†S. Martin and M. Vaughn, Phys. Rev. **D50**, 2282 (1994); Y. Yamada, Phys. Rev. **D50**, 3537 (1994); I. Jack, D.R.T. Jones, Phys. Lett. **B333**, 372 (1994)

J. Bagger, K. Matchev, D. Pierce, and R. Zhang, Nucl. Phys. **B491, 3 (1997)

- Calculate SUSY masses at the electroweak scale. Attach experimental errors to the masses.

- Calculation of production cross sections for $\tilde{t}_{1,2}$, $\tilde{b}_{1,2}$, $\tilde{\tau}_{1,2}$ with polarized e^- and e^+ beams.

- Fit the masses and production cross sections within given experimental errors by varying the low energy parameters: M_{E_k}, M_{L_k}, M_{D_k}, M_{Q_k}, M_{U_k}, A_τ, A_b, A_t, M_{H_1}, M_{H_2}, M_i, $\tan\beta$, $k = 1, 3$, $i = 1, 2, 3$

- use again RGEs to get high scale parameters within the errors.

This is a pure bottom-up approach!

mSUGRA

$\tan\beta = 10,\ M_0 = 200$ GeV, $M_{1/2} = 250$ GeV,
$A_0 = -100,\ \text{sign}(\mu) = 1$



1 σ error bands

GMSB

$M_M = 200$ TeV, $\Lambda = 100$ TeV, $N_5 = 1$, $\tan\beta = 15$, $A_0 = 0$, $\mathrm{sign}(\mu) = 1$



1 σ error bands

String

$\tan\beta = 10$, $M_{3/2} = 180$ GeV, $\sin^2\theta = 0.9$, O–I, $n_Q = 0$, $n_D = 1$, $n_U = -2$, $n_L = -3$, $n_E = -1$, and $n_{H_1} = n_{H_2} = -1$, sign$(\mu) = -1$







1 σ error bands

String Parameter Determination

First preliminary results

$m_{3/2}$	180	179.9 $\pm$ 0.1
t	14	13.6 $\pm$ 0.2
$g_s^2 \simeq 1/ < s >$	0.5	0.47 $\pm$ 0.04
δ_{GS}	0	-0.06 $\pm$ 0.03
$\tan\beta$	10	10 $\pm$ 0.1
n_{H_1}	-1	-1.06 $\pm$ 0.02
n_{H_2}	-1	-1.007$\pm$ 0.01
n_L	-3	-3.03$\pm$ 0.02
n_E	-1	-0.97$\pm$ 0.01
n_Q	0	-0.02$\pm$ 0.01
n_U	-2	-1.99$\pm$ 0.02
n_D	1	1.16$\pm$ 0.02

Summary

- We have used the bottom-up approach for determining the high scale parameters assuming we know masses and cross sections within the experimental errors.

- The model–independent reconstruction of the fundamental supersymmetric theory at the high scale, the grand unification scale M_{GUT} in supergravity or string scenarios, as well as the intermediate scale M_M in gauge mediated supersymmetry breaking, appears feasible.

- One can determine High Scale Parameters rather accurately. We have demonstrated this in case of GMSB as well as string theories.

- One has to use the most precise data available $\Rightarrow$ Combination of data from LHC and LC are very desirable if not crucial.