

Introduction to particle physics

Lecture 10: Quark Model

Frank Krauss

IPPP Durham

U Durham, Epiphany term 2010

Outline

- 1 Isospin
- 2 Adding (iso-)spins
- 3 Strange particles
- 4 The quark model

Isospin

Introduction: Discovery of neutrons

- Rutherford's experiment: Lightest atom = H (p - e -bound state)
But: next lightest atom (He) four times as heavy as hydrogen, with only two electrons. Similar for Li (three electrons, seven times as heavy), etc.. Why so heavy?
- Answer: Nuclei are bound states of protons and neutrons.
- Discovery of the Neutron by J.Chadwick (1932)
Bombard Beryllium with α -particles, very penetrating non-ionising radiation emerges. Send through paraffin, in turn protons are emitted. Measure speed of protons: original radiation cannot be γ 's.

Therefore new particle ("neutron") with nearly the same mass as the proton but no charge.

- Heisenberg's proposal (1932):
Neutron and proton are two manifestations of same state, **Nucleon**.

$$|\text{Nucleon}\rangle = \begin{vmatrix} \text{proton} \\ \text{neutron} \end{vmatrix} \rangle \text{ or } |N\rangle = \begin{vmatrix} p \\ n \end{vmatrix} \rangle$$

- Symmetry relating them: **Isospin** (very similar to spin).


(See also R. Myer's course)

Proposing mesons

- H.Yukawa (1934): First prediction of **mesons**.
- Answer to why neutrons and protons bind together in nucleus.
- Yukawa's underlying assumption: Introduce a new force, short-ranged, thus mediated by massive mesons.
- Estimate: 3-400 times the electron mass.
From uncertainty principle $\Delta E \Delta t \geq 1$ with time given by nucleon radius as $\Delta t \approx 1/r_0$. Assume r_0 of order $\mathcal{O}(1\text{fm})$, then $\Delta E \approx m_{\text{meson}} \approx 200 \text{ MeV}$
(Note: natural units used in this estimate).

The first “mesons”: The muon & the pion

- Two groups (1937): Anderson & Neddermeyer, Street & Stevenson:
Finding such particles in cosmic rays using cloud chambers.
- But: wrong lifetime (too long, indicating weaker interaction), and inconsistent mass measurements
- Two decisive experiments to clarify the situation (Rome, 1946 & Powell et al. in Bristol, 1947) with photo emulsions.
- Result: In fact two new particles:
 - One **weakly interacting, the muon**, μ (a lepton),
 - one **strongly interacting, the pion**, π (a meson).
Three differently charged versions, π^+ , π^- , π^0 ;
main decays: charged ones into muons plus a neutrino, neutral one into two γ 's.



Mesons and baryons

- Have two kinds of strongly interacting particles:
 - **Mesons** = bound states of a quark and an antiquark
 - **Baryons** = bound states of a three quarks or 3 antiquarks
- Quarks a point-like spin-1/2 particles
- **Quarks and gluons always in bound states** (strong interaction!)
- To accommodate for isospin: two quark-types, u and d :

$$\left| \begin{array}{c} u^{\text{P}} \\ d^{\text{DOWN}} \end{array} \right\rangle \text{ with isospin and third component } I_{u,d} = \frac{1}{2}, I_{u,d}^3 = \pm \frac{1}{2}.$$
- Then: $|p\rangle = |uud\rangle$, $|n\rangle = |udd\rangle$, $|\bar{p}\rangle = |\bar{u}\bar{u}\bar{d}\rangle$, $|\bar{n}\rangle = |\bar{u}\bar{d}\bar{d}\rangle$,
and $|\pi^+\rangle = |u\bar{d}\rangle$, $|\pi^-\rangle = |d\bar{u}\rangle$, $|\pi^0\rangle = |\bar{\pi}^0\rangle = \frac{1}{\sqrt{2}} [|u\bar{u}\rangle - |d\bar{d}\rangle]$.
- This also fixes the charges: $Q_u = \frac{2}{3}$, $Q_d = -\frac{1}{3}$.
- Obvious question: Are there more states?
 - yes, can have $|uuu\rangle$ and similar
 - yes, can supply the quarks with relative angular momentum etc..

Detour: (Iso-)Spins and their addition

Spin-1/2 systems: General remarks

- Spin-1/2 systems are often studied in physics.
- Spin-statistics theorem suggests that such systems are fermionic in nature, i.e. respect Pauli exclusion.
- Interesting in the context of this lecture:
Basic building blocks of matter (quarks & leptons) are spin-1/2.

- Simple representation:

$$|\uparrow\rangle = \left| \frac{1}{2}, +\frac{1}{2} \right\rangle \text{ and } |\downarrow\rangle = \left| \frac{1}{2}, -\frac{1}{2} \right\rangle.$$

Important: Distinguish total spin s and its projection, $s_z = s^3$ on a measurement axis (here the z -axis, could also be l and l^3).

- Examples: electron and its spin, isospin,
- Note: Spin can also occur as spin-1 etc..

Isospin assignments for the hadrons above:

$$|p, n\rangle = \left| \frac{1}{2}, \pm \frac{1}{2} \right\rangle \text{ and } |\pi^+, \pi^0, \pi^-\rangle = \left| \frac{1}{2}, \{1, 0, -1\} \right\rangle$$

Adding two spin-1/2 objects

- Often two spin-1/2 objects form a compound.
Examples: bound states of fermions, spin-orbit coupling, etc..
- If two spin-1/2 systems are added, the following objects can emerge:

$$|\uparrow\uparrow\rangle, |\uparrow\downarrow\rangle, |\downarrow\uparrow\rangle, \text{ and } |\downarrow\downarrow\rangle.$$

Naively, they have spin 1, 0, or -1, respectively.

But: **Need to distinguish total spin s and its projection onto the measurement axis s_z** (here, z has been chosen for simplicity)

- Then, truly relevant states are $s = 1$ (triplet, symmetric)

$$|1, 1\rangle = |\uparrow\uparrow\rangle, \quad |1, 0\rangle = \frac{1}{\sqrt{2}} [|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle], \quad |1, -1\rangle = |\downarrow\downarrow\rangle$$

and $s = 0$ (singlet, anti-symmetric):

$$|0, 0\rangle = \frac{1}{\sqrt{2}} [|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle]$$

- Catchy way of writing this: $\mathbf{2} \otimes \mathbf{2} = \mathbf{3} \oplus \mathbf{1}$

Clebsch-Gordan coefficients

- The **Clebsch-Gordan coefficients** in front of the new compound states can be calculated (or looked up).
- Formally speaking, they are defined as follows:

$$\left\langle s^{(1)}, s_z^{(1)}; s^{(2)}, s_z^{(2)} \mid s^{(1)}, s^{(2)}; s, s_z \right\rangle$$

indicating that two spin systems $s^{(1)}$ and $s^{(2)}$ are added to form a new spin system with total spin s (or J). Obviously, it is not only the total spin of each system that counts here, but also its orientation. This is typically indicated through “magnetic” quantum numbers, m , replacing the s_z in the literature.

Notation:	J	J	...
	M	M	...
m_1	m_2		
m_1	m_2	Coefficients	
.	.		
.	.		
.	.		

$1/2 \times 1/2$

			1	
		+1	1	0
+1/2	+1/2	1	0	0
+1/2	-1/2	1/2	1/2	1
-1/2	+1/2	1/2	-1/2	-1
		-1/2	-1/2	1

$1 \times 1/2$

				3/2	
			+3/2	3/2	1/2
+1	+1/2	1	-1/2	+1/2	
+1	-1/2	1/3	2/3	3/2	1/2
0	+1/2	2/3	-1/3	-1/2	-1/2
		0	-1/2	2/3	1/3
		-1	+1/2	1/3	-2/3
				-1	-1/2
					1

Note: Square-roots around the coefficients are understood in the table above

Using spin-algebra

- Identify: $|p\rangle = |\frac{1}{2}, +\frac{1}{2}\rangle$ and $|n\rangle = |\frac{1}{2}, -\frac{1}{2}\rangle$.
Heisenberg's proposal: Call this isospin (rather than spin).
- Also, the three kinds of pions can be written as:
 $|\pi^+\rangle = |1, +1\rangle$, $|\pi^0\rangle = |1, 0\rangle$, and $|\pi^-\rangle = |1, -1\rangle$.
- Catch: **Isospin conserved in strong interactions!**
- Dynamical implications: Bound states (here the deuteron).
Add two nucleons: can in principle have iso-singlet and iso-triplet.
But: No pp , nn -bound states, therefore $|d\rangle = |0, 0\rangle$ (deuteron = iso-singlet).
- Consider processes (+ their isospin amplitudes, below):

$$p + p \rightarrow d + \pi^+ \\ \mathcal{A}_{\text{iso}} \propto 1$$



$$p + n \rightarrow d + \pi^0 \\ \mathcal{A}_{\text{iso}} \propto 1/\sqrt{2}$$

$$n + n \rightarrow d + \pi^- \\ \mathcal{A}_{\text{iso}} \propto 1$$

Strangeness. Who ordered that?


Finding strange particles

- Rochester & Butler (1947): Cloud chamber experiment with cosmic rays. Unusual “fork” of a π^+ and a π^- .
- Interpretation: Cosmic ray particles, mass between π and p , the **kaon, K**.
- Like pions, but strangely long lifetime (typically decay to pions or a muon-neutrino pair), again hinting at weak interactions being responsible.
- Ultimately, this led to extending isospin ($SU(2)$) to a larger symmetry group, $SU(3)$ to catalogue mesons.



Finding more strangeness

- Anderson (1950): Another “strange” particle, decaying into proton and π^- , the **hyperon**: $\Lambda \rightarrow p\pi^-$.



- Copious production of such particles \rightarrow strong interaction
- But: Slow decays \rightarrow weak interaction

Cataloguing strangeness

- A new quantum number (like isospin)

(proposed in 1953 by Gell-Mann and Nishijima)

- Emerge as another quark type, strange, s :
 $I_s = 0$, $Q_s = Q_d = -\frac{1}{3}$, $S_s = -1$ ($S_{u,d} = 0$)
- **Conserved in strong interactions** (only produced pairwise, $s\bar{s}$),
 violated in weak interactions (s decays into u)
- Gell-Mann-Nishijima relation:

$$Q = e \left(I_3 + \frac{B+S}{2} \right).$$

(I_3 is the third component ($= \pm 1/2$ for p , n) of the isospin, $S = \pm 1$ for kaons, Λ 's, and Σ 's, B is the baryon number ($= 1$ for baryons like p , n , Λ , Σ and $= 0$ for mesons like π , K).

- Four mesons: kaons (K^+ , K^- , K^0 , \bar{K}^0).
 - All are pseudo-scalars (i.e. spin-0, negative parity), just like pions.
 - All have the same mass, about three times m_π ($m_\pi \approx 140$ MeV, $m_K \approx 495$ MeV) \implies "relatives"?

(But: "Funny" multiplet structure.)

The quark model


Quarks

- In 1964 Gell-Mann and Zweig proposed three “hypothetical” quarks, *u*p, *d*own and *s*trange, a mnemonic help.
All known hadrons are their “bound states”:
 - **Mesons** are made from a $q\bar{q}$ -pair, $q\bar{q} \equiv \mathbf{3} \otimes \bar{\mathbf{3}} = \mathbf{1} \oplus \mathbf{8}$
 - **baryons** from three quarks, $qqq \equiv \mathbf{3} \otimes \mathbf{3} \otimes \mathbf{3} = \mathbf{1} \oplus \mathbf{8} \oplus \mathbf{8} \oplus \mathbf{10}$;
- i.e. one singlet of mesons and baryons, one octet of mesons, and two octets and one decuplet of baryons.
- By now we know quarks are real (see next lecture).
 - Later more quark types were found (*c*harm, *b*ottom and *t*op).


(*u*, *d*, and *s* have masses around or below $\Lambda_{\text{QCD}} \approx 250$ MeV, the scale at which the strong coupling α_s diverges, and bound states emerge, *c*, *b*, and *t* are far above this scale.)

Meson multiplets

Pseudoscalars



Vectors



- Quark content of the mesons:


$$|K^+\rangle = |u\bar{s}\rangle, \quad |K^0\rangle = |d\bar{s}\rangle, \quad |K^-\rangle = |s\bar{u}\rangle, \quad |\bar{K}^0\rangle = |s\bar{d}\rangle,$$

$$|\pi^+\rangle = |u\bar{d}\rangle, \quad |\pi^0\rangle = \frac{1}{\sqrt{2}}|u\bar{u} - d\bar{d}\rangle, \quad |\pi^-\rangle = |d\bar{u}\rangle,$$


$$|\eta'\rangle = \frac{1}{\sqrt{6}}|u\bar{u} + d\bar{d} - 2s\bar{s}\rangle, \quad |\eta\rangle = \frac{1}{\sqrt{3}}|u\bar{u} + d\bar{d} + s\bar{s}\rangle.$$

Baryon multiplets

Octet




Decuplet




The discovery of the Ω^-

- In 1961, “tip” of the decuplet not yet found.
M.Gell-Mann’s prediction: $m = 1672$ MeV, plus the right production mechanism and a long lifetime.


- Decay chain:




(strangeness conserving)




($\Delta S = 1$ weak decay)



($\Delta S = 1$ weak decay)



($\Delta S = 0$ strong decay)



The postulate of colour

- In the decuplet, one problem appears: Some states like for instance the Δ^{++} are composed from three identical quarks (u 's for the Δ^{++}). Since the decuplet baryons are spin-3/2 objects they are fermions, i.e. their wave function must be antisymmetric. With three identical quarks, in identical spin states (spin-3/2 implies the spin-1/2's point into the same direction), this is possible only by invoking a new quantum number, **colour**.

We will discuss this when we encounter the strong interaction again.

Summary

- More particles in the zoo.
- First encounter with isospin as a first symmetry.
- Emergence of strangeness - giving rise to the quark model: $SU(3)$ or “the eightfold way”.
- Symmetry as **the** method of choice to gain control.
- Resonances as intermediate states.
- To read: Coughlan, Dodd & Gripaos, “The ideas of particle physics”, Sec 7-10.