

Exclusive photoproduction of vector mesons and Z/γ^*

Graeme Watt

University College London

*London workshop on Standard Model
discoveries with early LHC data*


30th March 2009

Talk based on work done in collaboration with L. Motyka:
Phys. Rev. D **78** (2008) 014023 [[arXiv:0805.2113](https://arxiv.org/abs/0805.2113)]

Motivation: central exclusive production at the LHC



Khoze, Martin, Ryskin [[arXiv:0802.0177](https://arxiv.org/abs/0802.0177), and references therein]:

$$\sigma(pp \rightarrow p+A+p) \sim \frac{S^2}{B_D^2} \left| \frac{\pi}{8} \int \frac{dQ_T^2}{Q_T^4} f_g(x_1, x'_1, Q_T^2, \mu^2) f_g(x_2, x'_2, Q_T^2, \mu^2) \right|^2 \hat{\sigma}(gg \rightarrow A)$$




$$f_g(x, x', Q_T^2, \mu^2) = R_g \frac{\partial}{\partial \ln Q_T^2} \left[\sqrt{T_g(Q_T^2, \mu^2)} x g(x, Q_T^2) \right]$$

Integrand
dominated by
 $Q_T \sim 1-2$ GeV
 \Rightarrow pQCD
 applicable (just).



Exclusive photoproduction at the LHC



Klein and Nystrand [[hep-ph/0311164](#)]


- Exclusive final state $E = J/\psi, \Upsilon, Z/\gamma^*$ with rapidity y .
- Flux dn/dk of quasi-real photons with energy $k \simeq (M_E/2) \exp(y) \simeq W^2/(2\sqrt{s})$

$$\frac{d\sigma}{dy}(h_1 h_2 \rightarrow h_1 + E + h_2) = k \frac{dn}{dk} \sigma(\gamma p \rightarrow E + p) \quad + \quad (y \rightarrow -y)$$

Disclaimer

- Neglect interference between photon–Pomeron and Pomeron–photon fusion, and effect of absorptive corrections from soft rescattering.
- Only present cross sections integrated over final state momenta, then these effects will be largely washed out. (Rapidity gap survival factor $S^2 \sim 0.7\text{--}0.9$.)
- Detailed treatment of these effects by Khoze, Martin and Ryskin [[hep-ph/0201301](#)] and by Rybarska, Schäfer and Szczurek [[arXiv:0705.2887](#), [arXiv:0805.0717](#)].


Exclusive photoproduction at HERA



- In LLA [Ryskin '93]:


$$\sigma \propto [xg(x, M_V^2/4)]^2$$

- Beyond LLA, use k_T -factorization [Martin, Ryskin, Teubner].




Small- x gluon density from exclusive J/ψ at HERA

Plot by P. Fleischmann (H1).




- J/ψ data can discriminate.
- ⇒ Fit gluon directly.

MRT+Nockles [[arXiv:0709.4406](https://arxiv.org/abs/0709.4406)]



Potential for odderon discovery at the Tevatron and LHC




- **Pomeron** is colour singlet exchange with **even** charge parity.
- Hypothetical partner with **odd** charge parity is the **odderon**.
- Bdzak, Motyka, Szymanowski, Cudell [[hep-ph/0702134](#)] calculated odderon-to-photon ratio using k_T -factorisation:

	J/ψ	Υ
Tevatron	0.3–0.6	0.8–1.7
LHC	0.06–0.15	0.16–0.38

- Odderon exchange also leads to different p_T -distribution.


Dipole model approach for exclusive diffractive processes

Bartels, Golec-Biernat, Peters [hep-ph/0301192]




- Non-forward photon impact factor calculated in the high-energy limit.

- Fourier transform from momentum space to coordinate space ($\mathbf{k} \rightarrow \mathbf{r}$), then to impact parameter space ($\Delta \rightarrow \mathbf{b}$), with $t = -\Delta^2$.



- Amplitude factorises into:
(wave function)·(dipole cross section)·(wave function).

Impact parameter dependent saturation (b-Sat) model



Kowalski, Teaney [[hep-ph/0304189](#)]

Kowalski, Motyka, G.W. [[hep-ph/0606272](#)]



G.W., Kowalski [[arXiv:0712.2670](#)]

- DGLAP-evolved gluon density with Gaussian b dependence:

$$\mathcal{N}(x, r, b) = 1 - \exp\left(-\frac{\pi^2}{2N_c} r^2 \alpha_s(\mu^2) x g(x, \mu^2) T(b)\right)$$


$$x g(x, \mu_0^2) = A_g x^{-\lambda_g} (1-x)^{5.6}, \quad T(b) = \frac{1}{2\pi B_G} e^{-\frac{b^2}{2B_G}}$$

- $B_G = 4 \text{ GeV}^{-2}$ from t -slope of exclusive J/ψ photoproduction.
- Fit input gluon parameters (μ_0^2, A_g, λ_g) to small- x F_2 data.
- Good (parameter-free) description of exclusive $\gamma^* p \rightarrow V + p$ ($V = \rho, \phi, J/\psi, \gamma$) and inclusive $F_2^{c\bar{c}}, F_2^{b\bar{b}}, F_L, F_2^D$.

Exclusive J/ψ photoproduction at Tevatron and LHC

Exclusive charmonium production [arXiv:0902.1271]

- CDF measure J/ψ :





$$\left. \frac{d\sigma}{dy} \right|_{y=0} = (3.92 \pm 0.62) \text{ nb.}$$

- Photoproduction contrib.: $3.4 \text{ nb} \times (S^2 \simeq 0.9)$
- Odderon-to-photon ratio: 0.3–0.6
- \Rightarrow Total theory prediction: (4.0–4.9) nb.

Prospects for J/ψ measurements at the LHC



ATLAS/CMS unlikely to measure exclusive J/ψ due to lack of low p_T trigger on leptons, but should be possible with ALICE/LHCb.

Exclusive Υ photoproduction at Tevatron and LHC




- Uncertainty in m_b and Ψ_γ
⇒ Scale to HERA data.
- Sensitive to W dependence of γp cross section.

Update: publication of ZEUS data [arXiv:0903.4205]



- ZEUS 2009 points move down cf. 2007 (prel.) points.
- Fit to HERA data more compatible with model.


Other theory predictions compared to HERA Υ data[ZEUS, [arXiv:0903.4205](https://arxiv.org/abs/0903.4205)]

- **RSS** = Rybarska, Schäfer, Szczurek [[arXiv:0805.0717](https://arxiv.org/abs/0805.0717)]
- **IKS** = Ivanov, Krasnikov, Szymanowski [[hep-ph/0412235](https://arxiv.org/abs/hep-ph/0412235)]
- **FMS** = Frankfurt, McDermott, Strikman [[hep-ph/9812316](https://arxiv.org/abs/hep-ph/9812316)]
- **MNRT** = Martin, Nockles, Ryskin, Teubner [[arXiv:0709.4406](https://arxiv.org/abs/0709.4406)]

Prospects for exclusive Υ at Tevatron and LHC

Tevatron


- Candidate events found by CDF.



- Cross section measurements eagerly awaited.


LHC

- Study by CMS [CMS PAS DIF-07-001] for 100 pb^{-1} and using STARLIGHT to generate signal sample.





Deeply virtual Compton scattering (DVCS) at HERA

- DVCS theoretically cleaner than exclusive vector meson production since no uncertainty from wave function.



Timelike Compton scattering (TCS) and $\gamma p \rightarrow Z + p$

- Analogous processes to DVCS at hadron–hadron colliders are TCS ($\gamma p \rightarrow \gamma^* + p$) and exclusive Z photoproduction.
- Wave functions for an incoming Z/γ^* with **spacelike** virtuality $q^2 = -Q^2 < 0$ are already known.
- Wave functions for an outgoing Z/γ^* with **timelike** $q^2 = M^2 > 0$ worked out by Motyka and G.W. [[arXiv:0805.2113](#)].
- Amplitude for $\gamma p \rightarrow \gamma^* + p$ is **not** simply the DVCS amplitude at $Q^2 = M^2$: pick up **real** contribution to the amplitude.
- TCS interferes at amplitude level with the pure QED subprocess ($\gamma\gamma \rightarrow \ell^+\ell^-$), but the latter can be precisely calculated and suppressed by a cut on the polar angle [Pire, Szymanowski, Wagner, [arXiv:0811.0321](#)]. (This is similar to the measurement of DVCS at HERA, which interferes with the Bethe–Heitler process.)

Exclusive Z photoproduction at Tevatron and LHC

Search for exclusive Z by CDF [arXiv:0902.2816]

- Eight candidate events found in 2.20 (2.03) fb^{-1} of data in the electron (muon) channel with $M_{\ell\ell} > 40 \text{ GeV}$ and $\eta_\ell < 4$, consistent with prediction for $\gamma\gamma \rightarrow \ell^+\ell^-$ (LPAIR).
- No candidate events in Z mass window \Rightarrow upper limit placed for exclusive Z cross section of $\sigma < 0.96 \text{ pb}$ at 95% confidence-level.
- Theory prediction: 0.3 fb , i.e. 3000 times lower than experimental limit.
- Certain BSM theory [A. White, hep-ph/0412062] predicts a much larger cross section, but without a quantitative estimate.
- Slightly more promising SM prediction at LHC: $\sigma = 13 \text{ fb}$.

Summary of predictions for J/ψ , Υ and Z^0 production

J/ψ	$d\sigma/dy _{y=0}$ (nb)	σ (nb)	Event rate (s^{-1})
Tevatron	3.4	28	0.33
LHC	9.8	120	71

$\Upsilon(1S)$	$d\sigma/dy _{y=0}$ (pb)	σ (pb)	Event rate (hr^{-1})
Tevatron	10	83	1.5
LHC	53	772	690

Z^0	$d\sigma/dy _{y=0}$ (fb)	σ (fb)	Event rate (yr^{-1})
Tevatron	0.077	0.30	0.065
LHC	1.4	13	135

- Event rates include leptonic branching ratio and assume a luminosity $\mathcal{L} = 2 \times 10^{32} \text{ cm}^{-2} \text{ s}^{-1}$ (Tevatron) and $\mathcal{L} = 10^{34} \text{ cm}^{-2} \text{ s}^{-1}$ (LHC). No gap survival factor included.